


	Number of wells analyzed (# of samples)	Number of wells w/ detects (# of samples)	Number of wells w/ detects located in a 'moderate' or 'high' vulnerability*	Number of wells w/ detects in areas of dense irrigation (>= 108 irrigation wells/township)*	Percentage of wells w/detects meeting either the vulnerability or well density criteria	Percentage of all wells meeting either the vulnerability or well density criteria	Number of wells exceeding 1/2 of respective MCL or other standard (and meeting vulnerability or well density criteria)	Number of wells exceeding respective MCL or other standard (and meeting vulnerability or well density criteria)
alachlor	141 (155)	1 (1)	0	0	0	62	0 (0)	0 (0)
atrazine	141 (155)	15 (18)	14	0	93	62	1 (1)	0 (0)
desethyl atrazine	64 (78)	13 (21)	13	0	100	73		
desisopropyl atrazine	64 (78)	2 (3)	2	0	100	73		
didealkyl atrazine	0	0	0	0	0	0		
metolachlor	141 (155)	0	0	0	0	62	0 (0)	0 (0)
simazine	53 (56)	4 (5)	4	0	100	68	0 (0)	0 (0)
Totals	141 (677)	23 (48)	21	0	91	62		

Percentage of Samples by Well Type (1990 - present)



Distribution of Concentrations from Samples with Detects




Figure 29. Summary of the samples analyzed for PMP herbicides since 1990 in the North Platte NRD. Taken from the Quality-assessed Agricultural Contaminant Database for Nebraska Ground Water, February, 2001 update. * see the vulnerability discussion for an explanation of these criteria.